

6. First Order Logic – Natural Deduction – Exercises

David Pereira José Proença Eduardo Tovar

RAMDE 2021/2022

Requirements and Model-driven Engineering

CISTER – ISEP

Porto, Portugal

<https://cister-labs.github.io/ramde2122>

Recalling Natural Deduction Rules in First Order Logic

Which are the new rules (on top of Propositional Logic)?

Elimination rule for \forall

If we know that $\forall x, \varphi$ holds, then we can conclude that φ holds for a specific term t

$$\frac{\forall x \varphi}{\varphi[t/x]} \forall E$$

Introduction rule for \forall

If we assume some term t and we are able to prove that $\varphi[t/x]$ then we can conclude that $\forall x, \varphi$.

$$\frac{[t] \quad \vdots \quad \varphi[t/x]}{\forall x \varphi[t/x]} \forall I$$

Which are the new rules (on top of Propositional Logic)?

Elimination rule for \exists

If we know that $\exists x, \varphi$ holds, and if assuming term t and $\varphi[t/x]$ we can deduce ψ , then we can prove ψ overall.

$$\frac{\begin{array}{c} [t \ \varphi[t/x]] \\ \vdots \\ \psi \end{array}}{\exists x \varphi} \exists E$$

Introduction rule for \exists

If we assume some term t and we are able to prove that $\varphi[t/x]$ then we can conclude that $\exists x, \varphi$.

$$\frac{\varphi[t/x]}{\exists x, \varphi} \exists I$$

Which are the new rules (on top of Propositional Logic)?

Elimination rule for =

If we know that two terms t_1 and t_2 are equal and that $\varphi[t_1/x]$ holds, then $\varphi[t_2/x]$ must also hold.

$$\frac{t_1 = t_2 \quad \varphi[t_1/x]}{\varphi[t_2/x]} =\mathbf{E}$$

Introduction rule for =

If we assume some term t and we are able to prove that $\varphi[t/x]$ then we can conclude that $\forall x, \varphi$.

$$\frac{}{t = t} =\mathbf{I}$$

Practical Exercises

Ex1. Build a proof of $\forall x(R(x) \wedge Q(x)) \vdash \forall xR(x) \wedge \forall xQ(x)$

Ex1. Build a proof of $\forall x(R(x) \wedge Q(x)) \vdash \forall xR(x) \wedge \forall xQ(x)$

$$1 \quad \underline{\forall x(R(x) \wedge Q(x))}$$

Ex1. Build a proof of $\forall x(R(x) \wedge Q(x)) \vdash \forall xR(x) \wedge \forall xQ(x)$

$$\begin{array}{l|l} 1 & \forall x(R(x) \wedge Q(x)) \\ \hline 2 & v \mid R(v) \wedge Q(v) \quad \forall E, 1 \end{array}$$

Ex1. Build a proof of $\forall x(R(x) \wedge Q(x)) \vdash \forall xR(x) \wedge \forall xQ(x)$

1		$\forall x(R(x) \wedge Q(x))$	
		<hr/>	
2		v $R(v) \wedge Q(v)$	$\forall E, 1$
3		$R(v)$	$\wedge E, 2$

Ex1. Build a proof of $\forall x(R(x) \wedge Q(x)) \vdash \forall xR(x) \wedge \forall xQ(x)$

1		$\forall x(R(x) \wedge Q(x))$	
<hr/>			
2		v $R(v) \wedge Q(v)$	$\forall E, 1$
3		$R(v)$	$\wedge E, 2$
4		$\forall xR(x)$	$\forall I, 2-3$

Ex1. Build a proof of $\forall x(R(x) \wedge Q(x)) \vdash \forall xR(x) \wedge \forall xQ(x)$

1		$\forall x(R(x) \wedge Q(x))$	
<hr/>			
2		$v \mid R(v) \wedge Q(v)$	$\forall E, 1$
3		$\mid R(v)$	$\wedge E, 2$
4		$\forall xR(x)$	$\forall I, 2-3$
5		$v \mid R(v) \wedge Q(v)$	$\forall E, 1$

Ex1. Build a proof of $\forall x(R(x) \wedge Q(x)) \vdash \forall xR(x) \wedge \forall xQ(x)$

1		$\forall x(R(x) \wedge Q(x))$	
<hr/>			
2		$R(v) \wedge Q(v)$	$\forall E, 1$
3		$R(v)$	$\wedge E, 2$
4		$\forall xR(x)$	$\forall I, 2-3$
5		$R(v) \wedge Q(v)$	$\forall E, 1$
6		$Q(v)$	$\wedge E, 5$

Ex1. Build a proof of $\forall x(R(x) \wedge Q(x)) \vdash \forall xR(x) \wedge \forall xQ(x)$

1		$\forall x(R(x) \wedge Q(x))$	
<hr/>			
2		$R(v) \wedge Q(v)$	$\forall E, 1$
3		$R(v)$	$\wedge E, 2$
4		$\forall xR(x)$	$\forall I, 2-3$
5		$R(v) \wedge Q(v)$	$\forall E, 1$
6		$Q(v)$	$\wedge E, 5$
7		$\forall xQ(x)$	$\forall I, 5-7$

Ex1. Build a proof of $\forall x(R(x) \wedge Q(x)) \vdash \forall xR(x) \wedge \forall xQ(x)$

1	$\forall x(R(x) \wedge Q(x))$	
<hr/>		
2	v $R(v) \wedge Q(v)$	$\forall E, 1$
3	$R(v)$	$\wedge E, 2$
4	$\forall xR(x)$	$\forall I, 2-3$
5	v $R(v) \wedge Q(v)$	$\forall E, 1$
6	$Q(v)$	$\wedge E, 5$
7	$\forall xQ(x)$	$\forall I, 5-7$
8	$\forall xR(x) \wedge \forall xQ(x)$	$\wedge I, 4, 8$

Ex2. Build a proof of $\forall x(R(x) \rightarrow Q(x)) \vdash \forall xR(x) \rightarrow \forall xQ(x)$

Ex2. Build a proof of $\forall x(R(x) \rightarrow Q(x)) \vdash \forall xR(x) \rightarrow \forall xQ(x)$

1 $\boxed{\forall x(R(x) \rightarrow Q(x))}$

Ex2. Build a proof of $\forall x(R(x) \rightarrow Q(x)) \vdash \forall xR(x) \rightarrow \forall xQ(x)$

1		$\forall x(R(x) \rightarrow Q(x))$

2		$\forall xR(x)$

Ex2. Build a proof of $\forall x(R(x) \rightarrow Q(x)) \vdash \forall xR(x) \rightarrow \forall xQ(x)$

1	$\forall x(R(x) \rightarrow Q(x))$	
2	$\forall xR(x)$	
3	$v \mid R(v) \rightarrow Q(v)$	$\forall E, 1$

Ex2. Build a proof of $\forall x(R(x) \rightarrow Q(x)) \vdash \forall xR(x) \rightarrow \forall xQ(x)$

1		$\forall x(R(x) \rightarrow Q(x))$	
		<hr/>	
2			
3			
4			

$R(v) \rightarrow Q(v)$ $\forall E, 1$

$R(v)$ $\forall E, 2$

Ex2. Build a proof of $\forall x(R(x) \rightarrow Q(x)) \vdash \forall xR(x) \rightarrow \forall xQ(x)$

1		$\forall x(R(x) \rightarrow Q(x))$	
<hr/>			
2			
		$\forall xR(x)$	
<hr/>			
3			
		v	
		$R(v) \rightarrow Q(v)$	$\forall E, 1$
4			
		$R(v)$	$\forall E, 2$
5			
		$Q(v)$	$\Rightarrow E, 3, 4$

Ex2. Build a proof of $\forall x(R(x) \rightarrow Q(x)) \vdash \forall xR(x) \rightarrow \forall xQ(x)$

1		$\forall x(R(x) \rightarrow Q(x))$			
<hr/>					
2			$\forall xR(x)$		
<hr/>					
3				$R(v) \rightarrow Q(v)$	$\forall E, 1$
4				$R(v)$	$\forall E, 2$
5				$Q(v)$	$\Rightarrow E, 3, 4$
6			$\forall xQ(x)$	$\forall I, 3-5$	

Ex2. Build a proof of $\forall x(R(x) \rightarrow Q(x)) \vdash \forall xR(x) \rightarrow \forall xQ(x)$

1	$\forall x(R(x) \rightarrow Q(x))$	
2	$\forall xR(x)$	
3	v $R(v) \rightarrow Q(v)$	$\forall E, 1$
4	$R(v)$	$\forall E, 2$
5	$Q(v)$	$\Rightarrow E, 3, 4$
6	$\forall xQ(x)$	$\forall I, 3-5$
7	$\forall xP(x) \rightarrow \forall xQ(x)$	$\Rightarrow I, 2-6$

Ex3. Build a proof of $\exists x(R(x) \rightarrow Q(x)) \vdash \exists xR(x) \rightarrow \exists xQ(x)$

Ex3. Build a proof of $\exists x(R(x) \rightarrow Q(x)) \vdash \exists xR(x) \rightarrow \exists xQ(x)$

$$1 \quad \underline{\exists x(R(x) \rightarrow Q(x))}$$

Ex3. Build a proof of $\exists x(R(x) \rightarrow Q(x)) \vdash \exists xR(x) \rightarrow \exists xQ(x)$

$$\begin{array}{l|l} 1 & \exists x(R(x) \rightarrow Q(x)) \\ \hline 2 & \begin{array}{|l} \exists xR(x) \end{array} \end{array}$$

Ex3. Build a proof of $\exists x(R(x) \rightarrow Q(x)) \vdash \exists xR(x) \rightarrow \exists xQ(x)$

$$\begin{array}{l|l} 1 & \exists x(R(x) \rightarrow Q(x)) \\ \hline 2 & \left| \begin{array}{l} \exists xR(x) \\ \hline 3 \quad \left| \begin{array}{l} v \quad R(v) \rightarrow Q(v) \\ \hline \end{array} \right. \end{array} \right. \end{array}$$

Ex3. Build a proof of $\exists x(R(x) \rightarrow Q(x)) \vdash \exists xR(x) \rightarrow \exists xQ(x)$

1		$\exists x(R(x) \rightarrow Q(x))$			
		<hr/>			
2			$\exists xR(x)$		
			<hr/>		
3				$R(v) \rightarrow Q(v)$	
				<hr/>	
4					$R(v)$
					<hr/>

Ex3. Build a proof of $\exists x(R(x) \rightarrow Q(x)) \vdash \exists xR(x) \rightarrow \exists xQ(x)$

1		$\exists x(R(x) \rightarrow Q(x))$	

2		$\exists xR(x)$	

3		v $R(v) \rightarrow Q(v)$	

4		$R(v)$	

5		$Q(v)$	$\Rightarrow E, 3, 4$

Ex3. Build a proof of $\exists x(R(x) \rightarrow Q(x)) \vdash \exists xR(x) \rightarrow \exists xQ(x)$

1		$\exists x(R(x) \rightarrow Q(x))$	
2			
3			
4			
5			$\Rightarrow E, 3, 4$
6			$\exists I, 5$

Ex3. Build a proof of $\exists x(R(x) \rightarrow Q(x)) \vdash \exists xR(x) \rightarrow \exists xQ(x)$

1		$\exists x(R(x) \rightarrow Q(x))$	
2			
3			
4			
5			$\Rightarrow E, 3, 4$
6			$\exists I, 5$
7			$\exists E, 2, 4-6$

Ex3. Build a proof of $\exists x(R(x) \rightarrow Q(x)) \vdash \exists xR(x) \rightarrow \exists xQ(x)$

1	$\exists x(R(x) \rightarrow Q(x))$ <hr style="border: 0.5px solid black; margin-top: 5px;"/>	
2	$\exists xR(x)$ <hr style="border: 0.5px solid black; margin-top: 5px;"/>	
3	v $R(v) \rightarrow Q(v)$ <hr style="border: 0.5px solid black; margin-top: 5px;"/>	
4	v $R(v)$ <hr style="border: 0.5px solid black; margin-top: 5px;"/>	
5	v $Q(v)$	$\Rightarrow E, 3, 4$
6	v $\exists xQ(x)$	$\exists I, 5$
7	$\exists xQ(x)$	$\exists E, 2, 4-6$
8	$\exists xQ(x)$	$\exists E, 1, 3-7$

Ex3. Build a proof of $\exists x(R(x) \rightarrow Q(x)) \vdash \exists xR(x) \rightarrow \exists xQ(x)$

1	$\exists x(R(x) \rightarrow Q(x))$	
2	$\exists xR(x)$	
3	v $R(v) \rightarrow Q(v)$	
4	$R(v)$	
5	$Q(v)$	$\Rightarrow E, 3, 4$
6	$\exists xQ(x)$	$\exists I, 5$
7	$\exists xQ(x)$	$\exists E, 2, 4-6$
8	$\exists xQ(x)$	$\exists E, 1, 3-7$
9	$\exists xP(x) \rightarrow \exists xQ(x)$	$\Rightarrow I, 2-8$

Ex4. Build a proof of $\exists x\neg Q(x) \vdash \neg\forall xQ(x)$

Ex4. Build a proof of $\exists x\neg Q(x) \vdash \neg\forall xQ(x)$

1 $\boxed{\exists x\neg Q(x)}$

Ex4. Build a proof of $\exists x\neg Q(x) \vdash \neg\forall xQ(x)$

$$\begin{array}{l|l} 1 & \exists x\neg Q(x) \\ \hline 2 & \boxed{\forall xQ(x)} \end{array}$$

Ex4. Build a proof of $\exists x\neg Q(x) \vdash \neg\forall xQ(x)$

$$\begin{array}{l|l|l} 1 & \exists x\neg Q(x) & \\ \hline 2 & \forall xQ(x) & \\ \hline 3 & v \mid \neg Q(v) & \end{array}$$

Ex4. Build a proof of $\exists x\neg Q(x) \vdash \neg\forall xQ(x)$

1		$\exists x\neg Q(x)$		

2			$\forall xQ(x)$	

3				$\neg Q(v)$

4				$Q(v)$
				$\forall E, 2$

Ex4. Build a proof of $\exists x\neg Q(x) \vdash \neg\forall xQ(x)$

1		$\exists x\neg Q(x)$			

2			$\forall xQ(x)$		

3				$\neg Q(v)$	

4				$Q(v)$	$\forall E, 2$
5				\perp	$\perp I, 3, 4$

Ex4. Build a proof of $\exists x\neg Q(x) \vdash \neg\forall xQ(x)$

1		$\exists x\neg Q(x)$	
<hr/>			
2		$\forall xQ(x)$	
<hr/>			
3		v $\neg Q(v)$	
<hr/>			
4		$Q(v)$	$\forall E, 2$
5		\perp	$\perp I, 3, 4$
6		\perp	$\exists E, 1, 3-5$

Ex4. Build a proof of $\exists x\neg Q(x) \vdash \neg\forall xQ(x)$

1	$\exists x\neg Q(x)$	
2	$\forall xQ(x)$	
3	$\neg Q(v)$	
4	$Q(v)$	$\forall E, 2$
5	\perp	$\perp I, 3, 4$
6	\perp	$\exists E, 1, 3-5$
7	$\neg\forall xQ(x)$	$\neg I, 2-6$

Ex5. Build a proof of $\neg\exists x\neg Q(x) \vdash \forall xQ(x)$

Ex5. Build a proof of $\neg\exists x\neg Q(x) \vdash \forall xQ(x)$

1 $\neg\exists x\neg Q(x)$

Ex5. Build a proof of $\neg\exists x\neg Q(x) \vdash \forall xQ(x)$

$$\begin{array}{l|l} 1 & \neg\exists x\neg Q(x) \\ \hline 2 & \boxed{\neg\forall xQ(x)} \end{array}$$

Ex5. Build a proof of $\neg\exists x\neg Q(x) \vdash \forall xQ(x)$

$$\begin{array}{l|l|l} 1 & \neg\exists x\neg Q(x) & \\ \hline 2 & \neg\forall xQ(x) & \\ \hline 3 & v & \neg Q(v) \end{array}$$

Ex5. Build a proof of $\neg\exists x\neg Q(x) \vdash \forall xQ(x)$

1		$\neg\exists x\neg Q(x)$			
2			$\neg\forall xQ(x)$		
3				$\neg Q(v)$	
4				$\exists x\neg Q(x)$	$\exists I, 3$

Ex5. Build a proof of $\neg\exists x\neg Q(x) \vdash \forall xQ(x)$

1		$\neg\exists x\neg Q(x)$		
2			$\neg\forall xQ(x)$	
3				$\neg Q(v)$
4				$\exists x\neg Q(x)$ $\exists I, 3$
5				\perp $\perp I, 1, 4$

Ex5. Build a proof of $\neg\exists x\neg Q(x) \vdash \forall xQ(x)$

1		$\neg\exists x\neg Q(x)$			
2			$\neg\forall xQ(x)$		
3				$\neg Q(v)$	
4				$\exists x\neg Q(x)$	$\exists I, 3$
5				\perp	$\perp I, 1, 4$
6				$\neg\neg Q(v)$	$\neg I, 3-5$

Ex5. Build a proof of $\neg\exists x\neg Q(x) \vdash \forall xQ(x)$

1		$\neg\exists x\neg Q(x)$			
2			$\neg\forall xQ(x)$		
3				$\neg Q(v)$	
4				$\exists x\neg Q(x)$	$\exists I, 3$
5				\perp	$\perp I, 1, 4$
6				$\neg\neg Q(v)$	$\neg I, 3-5$
7				$Q(v)$	$\neg E, 6$

Ex5. Build a proof of $\neg\exists x\neg Q(x) \vdash \forall xQ(x)$

1		$\neg\exists x\neg Q(x)$			
2			$\neg\forall xQ(x)$		
3				$\neg Q(v)$	
4				$\exists x\neg Q(x)$	$\exists I, 3$
5				\perp	$\perp I, 1, 4$
6				$\neg\neg Q(v)$	$\neg I, 3-5$
7				$Q(v)$	$\neg E, 6$
8				$\forall xQ(x)$	$\forall I, 3-7$

Ex5. Build a proof of $\neg\exists x\neg Q(x) \vdash \forall xQ(x)$

1		$\neg\exists x\neg Q(x)$			
2			$\neg\forall xQ(x)$		
3				$\neg Q(v)$	
4				$\exists x\neg Q(x)$	$\exists I, 3$
5				\perp	$\perp I, 1, 4$
6				$\neg\neg Q(v)$	$\neg I, 3-5$
7				$Q(v)$	$\neg E, 6$
8			$\forall xQ(x)$	$\forall I, 3-7$	
9		\perp	$\perp I, 2, 8$		

Ex5. Build a proof of $\neg\exists x\neg Q(x) \vdash \forall xQ(x)$

1	$\neg\exists x\neg Q(x)$		
2	$\neg\forall xQ(x)$		
3	v	$\neg Q(v)$	
4		$\exists x\neg Q(x)$	$\exists I, 3$
5		\perp	$\perp I, 1, 4$
6		$\neg\neg Q(v)$	$\neg I, 3-5$
7		$Q(v)$	$\neg E, 6$
8	$\forall xQ(x)$		$\forall I, 3-7$
9	\perp		$\perp I, 2, 8$
10	$\neg\neg\forall xQ(x)$		$\neg I, 2-9$

Ex5. Build a proof of $\neg\exists x\neg Q(x) \vdash \forall xQ(x)$

1	$\neg\exists x\neg Q(x)$		
2	$\neg\forall xQ(x)$		
3	v	$\neg Q(v)$	
4		$\exists x\neg Q(x)$	$\exists I, 3$
5		\perp	$\perp I, 1, 4$
6		$\neg\neg Q(v)$	$\neg I, 3-5$
7		$Q(v)$	$\neg E, 6$
8	$\forall xQ(x)$		$\forall I, 3-7$
9	\perp		$\perp I, 2, 8$
10	$\neg\neg\forall xQ(x)$		$\neg I, 2-9$
11	$\forall xQ(x)$		$\neg E, 10$

Ex6. Build a proof of $\forall xQ(x) \vdash \neg\exists x\neg Q(x)$

Ex6. Build a proof of $\forall xQ(x) \vdash \neg\exists x\neg Q(x)$

1 $\underbrace{\forall xQ(x)}$

Ex6. Build a proof of $\forall xQ(x) \vdash \neg\exists x\neg Q(x)$

1		$\forall xQ(x)$	

2			
			$\exists x\neg Q(x)$

Ex6. Build a proof of $\forall xQ(x) \vdash \neg\exists x\neg Q(x)$

1		$\forall xQ(x)$	

2			$\exists x\neg Q(x)$

3			$\neg\forall xQ(x)$
			Ex4., 1

Ex6. Build a proof of $\forall xQ(x) \vdash \neg\exists x\neg Q(x)$

1	$\forall xQ(x)$	
2	$\exists x\neg Q(x)$	
3	$\neg\forall xQ(x)$	Ex4., 1
4	\perp	\perp I, 1, 3

Ex6. Build a proof of $\forall xQ(x) \vdash \neg\exists x\neg Q(x)$

1	$\forall xQ(x)$	

2	$\exists x\neg Q(x)$	

3	$\neg\forall xQ(x)$	Ex4., 1
4	\perp	\perp I, 1, 3
5	$\neg\exists xQ(x)$	\neg I, 2-4

Ex7. Build a proof of $\exists x(P(x) \wedge Q(x)), \forall x(P(x) \rightarrow R(x)) \vdash \exists x(R(x) \wedge P(x))$

1 | $\exists x(P(x) \wedge Q(x))$

Ex7. Build a proof of $\exists x(P(x) \wedge Q(x)), \forall x(P(x) \rightarrow R(x)) \vdash \exists x(R(x) \wedge P(x))$

- 1 $\exists x(P(x) \wedge Q(x))$
- 2 $\forall x(P(x) \rightarrow R(x))$

Ex7. Build a proof of $\exists x(P(x) \wedge Q(x)), \forall x(P(x) \rightarrow R(x)) \vdash \exists x(R(x) \wedge P(x))$

1		$\exists x(P(x) \wedge Q(x))$
2		$\forall x(P(x) \rightarrow R(x))$
<hr/>		
3		v $P(v) \wedge Q(v)$

Ex7. Build a proof of $\exists x(P(x) \wedge Q(x)), \forall x(P(x) \rightarrow R(x)) \vdash \exists x(R(x) \wedge P(x))$

1		$\exists x(P(x) \wedge Q(x))$	
2		$\forall x(P(x) \rightarrow R(x))$	
		<hr/>	
3		v $P(v) \wedge Q(v)$	
		<hr/>	
4		$P(v)$	$\wedge E, 3$

Ex7. Build a proof of $\exists x(P(x) \wedge Q(x)), \forall x(P(x) \rightarrow R(x)) \vdash \exists x(R(x) \wedge P(x))$

1		$\exists x(P(x) \wedge Q(x))$	
2		$\forall x(P(x) \rightarrow R(x))$	
		<hr/>	
3		v	$P(v) \wedge Q(v)$
			<hr/>
4			$P(v)$ $\wedge E, 3$
5			$P(v) \rightarrow R(v)$ $\forall E, 2$

Ex7. Build a proof of $\exists x(P(x) \wedge Q(x)), \forall x(P(x) \rightarrow R(x)) \vdash \exists x(R(x) \wedge P(x))$

1		$\exists x(P(x) \wedge Q(x))$	
2		$\forall x(P(x) \rightarrow R(x))$	
		<hr/>	
3		v	$P(v) \wedge Q(v)$
			<hr/>
4			$P(v)$ $\wedge E, 3$
5			$P(v) \rightarrow R(v)$ $\forall E, 2$
6			$R(v)$ $\Rightarrow E, 4, 5$

Ex7. Build a proof of $\exists x(P(x) \wedge Q(x)), \forall x(P(x) \rightarrow R(x)) \vdash \exists x(R(x) \wedge P(x))$

1		$\exists x(P(x) \wedge Q(x))$	
2		$\forall x(P(x) \rightarrow R(x))$	
		<hr/>	
3		v	$P(v) \wedge Q(v)$
			<hr/>
4			$P(v)$ $\wedge E, 3$
5			$P(v) \rightarrow R(v)$ $\forall E, 2$
6			$R(v)$ $\Rightarrow E, 4, 5$
7			$Q(v)$ $\wedge E, 3$

Ex7. Build a proof of $\exists x(P(x) \wedge Q(x)), \forall x(P(x) \rightarrow R(x)) \vdash \exists x(R(x) \wedge P(x))$

1		$\exists x(P(x) \wedge Q(x))$	
2		$\forall x(P(x) \rightarrow R(x))$	
		<hr/>	
3		v $P(v) \wedge Q(v)$	
		<hr/>	
4		$P(v)$	$\wedge E, 3$
5		$P(v) \rightarrow R(v)$	$\forall E, 2$
6		$R(v)$	$\Rightarrow E, 4, 5$
7		$Q(v)$	$\wedge E, 3$
8		$R(v) \wedge Q(v)$	$\wedge I, 6, 7$

Ex7. Build a proof of $\exists x(P(x) \wedge Q(x)), \forall x(P(x) \rightarrow R(x)) \vdash \exists x(R(x) \wedge P(x))$

1		$\exists x(P(x) \wedge Q(x))$	
2		$\forall x(P(x) \rightarrow R(x))$	
		<hr/>	
3	v	$P(v) \wedge Q(v)$	
		<hr/>	
4		$P(v)$	$\wedge E, 3$
5		$P(v) \rightarrow R(v)$	$\forall E, 2$
6		$R(v)$	$\Rightarrow E, 4, 5$
7		$Q(v)$	$\wedge E, 3$
8		$R(v) \wedge Q(v)$	$\wedge I, 6, 7$
9		$\exists x(R(x) \wedge Q(x))$	$\exists I, 8$

Ex7. Build a proof of $\exists x(P(x) \wedge Q(x)), \forall x(P(x) \rightarrow R(x)) \vdash \exists x(R(x) \wedge P(x))$

1		$\exists x(P(x) \wedge Q(x))$	
2		$\forall x(P(x) \rightarrow R(x))$	
		<hr/>	
3	v	$P(v) \wedge Q(v)$	
		<hr/>	
4		$P(v)$	$\wedge E, 3$
5		$P(v) \rightarrow R(v)$	$\forall E, 2$
6		$R(v)$	$\Rightarrow E, 4, 5$
7		$Q(v)$	$\wedge E, 3$
8		$R(v) \wedge Q(v)$	$\wedge I, 6, 7$
9		$\exists x(R(x) \wedge Q(x))$	$\exists I, 8$
10		$\exists x(R(x) \wedge Q(x))$	$\exists E, 1, 3-9$

Ex8. Build a proof of $\forall x \forall y (x = y \rightarrow f(x) = f(y))$

Ex8. Build a proof of $\forall x \forall y (x = y \rightarrow f(x) = f(y))$

$$1 \quad | \quad u \quad | \quad v \quad | \quad \underline{u = v}$$

Ex8. Build a proof of $\forall x \forall y (x = y \rightarrow f(x) = f(y))$

$$\begin{array}{l} 1 \quad \left| \begin{array}{|l} u \\ v \end{array} \right| \quad \left| \begin{array}{l} u = v \\ \hline \neg(f(u) = f(v)) \end{array} \right. \end{array}$$

Ex8. Build a proof of $\forall x \forall y (x = y \rightarrow f(x) = f(y))$

$$\begin{array}{l|l|l|l} 1 & u & v & u = v \\ 2 & & & \neg(f(u) = f(v)) \\ 3 & & & \hline & & & f(u) = f(u) \quad =I \end{array}$$

Ex8. Build a proof of $\forall x \forall y (x = y \rightarrow f(x) = f(y))$

1		u		v		$u = v$	
2						$\neg(f(u) = f(v))$	
3						$f(u) = f(u)$	=I
4						$f(u) = f(v)$	=E, 1, 3

Ex8. Build a proof of $\forall x \forall y (x = y \rightarrow f(x) = f(y))$

1		u		v		$u = v$	
2						$\neg(f(u) = f(v))$	
3						$f(u) = f(u)$	=I
4						$f(u) = f(v)$	=E, 1, 3
5						\perp	\perp I, 2-4

Ex8. Build a proof of $\forall x \forall y (x = y \rightarrow f(x) = f(y))$

1		u		v		$u = v$	
2						$\neg(f(u) = f(v))$	
3						$f(u) = f(u)$	=I
4						$f(u) = f(v)$	=E, 1, 3
5						\perp	\perp I, 2-4
6						$\neg\neg(f(u) = f(v))$	\neg I, 2-5

Ex8. Build a proof of $\forall x \forall y (x = y \rightarrow f(x) = f(y))$

1		u		v		$u = v$	
2						$\neg(f(u) = f(v))$	
3						$f(u) = f(u)$	=I
4						$f(u) = f(v)$	=E, 1, 3
5						\perp	\perp I, 2-4
6						$\neg\neg(f(u) = f(v))$	\neg I, 2-5
7						$f(u) = f(v)$	\neg E, 6

Ex8. Build a proof of $\forall x \forall y (x = y \rightarrow f(x) = f(y))$

1	u	v	$u = v$		
2			$\neg(f(u) = f(v))$		
3			$f(u) = f(u)$	=I	
4			$f(u) = f(v)$	=E, 1, 3	
5			\perp	\perp I, 2-4	
6			$\neg\neg(f(u) = f(v))$	\neg I, 2-5	
7			$f(u) = f(v)$	\neg E, 6	
8			$u = v \rightarrow f(u) = f(v)$	\Rightarrow I, 1-7	

Ex8. Build a proof of $\forall x \forall y (x = y \rightarrow f(x) = f(y))$

1			$u = v$	
2			$\neg(f(u) = f(v))$	
3			$f(u) = f(u)$	=I
4			$f(u) = f(v)$	=E, 1, 3
5			\perp	\perp I, 2-4
6			$\neg\neg(f(u) = f(v))$	\neg I, 2-5
7			$f(u) = f(v)$	\neg E, 6
8		$u = v \rightarrow f(u) = f(v)$		\Rightarrow I, 1-7
9	$\forall y (u = y \rightarrow f(u) = f(y))$			\forall I, 8

Ex8. Build a proof of $\forall x \forall y (x = y \rightarrow f(x) = f(y))$

1	u	v	$u = v$	
2			$\neg(f(u) = f(v))$	
3			$f(u) = f(u)$	$=I$
4			$f(u) = f(v)$	$=E, 1, 3$
5			\perp	$\perp I, 2-4$
6			$\neg\neg(f(u) = f(v))$	$\neg I, 2-5$
7			$f(u) = f(v)$	$\neg E, 6$
8			$u = v \rightarrow f(u) = f(v)$	$\Rightarrow I, 1-7$
9			$\forall y (u = y \rightarrow f(u) = f(y))$	$\forall I, 8$
10			$\forall x \forall y (x = y \rightarrow f(x) = f(y))$	$\forall I, 9$

Ex9. Build a proof of $\forall x P(a, x, x), \forall x \forall y \forall z (P(x, y, z) \rightarrow P(f(x), y, f(z))) \vdash P(f(a), a, f(a))$

Ex9. Build a proof of $\forall xP(a, x, x), \forall x\forall y\forall z(P(x, y, z) \rightarrow P(f(x), y, f(z))) \vdash P(f(a), a, f(a))$

1 | $\forall xP(a, x, x)$

Ex9. Build a proof of $\forall xP(a, x, x), \forall x\forall y\forall z(P(x, y, z) \rightarrow P(f(x), y, f(z))) \vdash P(f(a), a, f(a))$

- 1 $\forall xP(a, x, x)$
- 2 $\forall x\forall y\forall z(P(x, y, z) \rightarrow P(f(x), y, f(z)))$

Ex9. Build a proof of $\forall xP(a, x, x), \forall x\forall y\forall z(P(x, y, z) \rightarrow P(f(x), y, f(z))) \vdash P(f(a), a, f(a))$

1	$\forall xP(a, x, x)$	
2	$\forall x\forall y\forall z(P(x, y, z) \rightarrow P(f(x), y, f(z)))$	
<hr/>		
3	$P(a, a, a)$	$\forall E, 1$

Ex9. Build a proof of $\forall xP(a, x, x), \forall x\forall y\forall z(P(x, y, z) \rightarrow P(f(x), y, f(z))) \vdash P(f(a), a, f(a))$

1	$\forall xP(a, x, x)$	
2	$\forall x\forall y\forall z(P(x, y, z) \rightarrow P(f(x), y, f(z)))$	
<hr/>		
3	$P(a, a, a)$	$\forall E, 1$
4	$\forall y\forall z(P(a, y, z) \rightarrow P(f(a), y, f(z)))$	$\forall E, 2$

Ex9. Build a proof of $\forall xP(a, x, x), \forall x\forall y\forall z(P(x, y, z) \rightarrow P(f(x), y, f(z))) \vdash P(f(a), a, f(a))$

1	$\forall xP(a, x, x)$	
2	$\forall x\forall y\forall z(P(x, y, z) \rightarrow P(f(x), y, f(z)))$	
<hr/>		
3	$P(a, a, a)$	$\forall E, 1$
4	$\forall y\forall z(P(a, y, z) \rightarrow P(f(a), y, f(z)))$	$\forall E, 2$
5	$\forall z(P(a, a, z) \rightarrow P(f(a), a, f(z)))$	$\forall E, 4$

Ex9. Build a proof of $\forall xP(a, x, x), \forall x\forall y\forall z(P(x, y, z) \rightarrow P(f(x), y, f(z))) \vdash P(f(a), a, f(a))$

1	$\forall xP(a, x, x)$	
2	$\forall x\forall y\forall z(P(x, y, z) \rightarrow P(f(x), y, f(z)))$	
<hr/>		
3	$P(a, a, a)$	$\forall E, 1$
4	$\forall y\forall z(P(a, y, z) \rightarrow P(f(a), y, f(z)))$	$\forall E, 2$
5	$\forall z(P(a, a, z) \rightarrow P(f(a), a, f(z)))$	$\forall E, 4$
6	$P(a, a, a) \rightarrow P(f(a), a, f(a))$	$\forall E, 5$

Ex9. Build a proof of $\forall xP(a, x, x), \forall x\forall y\forall z(P(x, y, z) \rightarrow P(f(x), y, f(z))) \vdash P(f(a), a, f(a))$

1	$\forall xP(a, x, x)$	
2	$\forall x\forall y\forall z(P(x, y, z) \rightarrow P(f(x), y, f(z)))$	
<hr/>		
3	$P(a, a, a)$	$\forall E, 1$
4	$\forall y\forall z(P(a, y, z) \rightarrow P(f(a), y, f(z)))$	$\forall E, 2$
5	$\forall z(P(a, a, z) \rightarrow P(f(a), a, f(z)))$	$\forall E, 4$
6	$P(a, a, a) \rightarrow P(f(a), a, f(a))$	$\forall E, 5$
7	$P(f(a), a, f(a))$	$\Rightarrow E, 3, 6$

Ex10. Build a proof of $\exists x\exists y(H(x, y) \vee H(y, x)), \neg\exists xH(x, x) \vdash \exists x\exists y\neg(x = y)$

1 | $\exists x\exists y(H(x, y) \vee H(y, x))$

Ex10. Build a proof of $\exists x\exists y(H(x, y) \vee H(y, x)), \neg\exists xH(x, x) \vdash \exists x\exists y\neg(x = y)$

- 1 $\exists x\exists y(H(x, y) \vee H(y, x))$
- 2 $\neg\exists xH(x, x)$

Ex10. Build a proof of $\exists x\exists y(H(x, y) \vee H(y, x)), \neg\exists xH(x, x) \vdash \exists x\exists y\neg(x = y)$

1		$\exists x\exists y(H(x, y) \vee H(y, x))$
2		$\neg\exists xH(x, x)$
<hr/>		
3		u v $u = v$

Ex10. Build a proof of $\exists x\exists y(H(x,y) \vee H(y,x)), \neg\exists xH(x,x) \vdash \exists x\exists y\neg(x = y)$

1		$\exists x\exists y(H(x,y) \vee H(y,x))$			
2		$\neg\exists xH(x,x)$			

3		u	v		$u = v$

4					$H(u,v) \vee H(v,u)$

5					$H(u,v)$

Ex10. Build a proof of $\exists x\exists y(H(x,y) \vee H(y,x)), \neg\exists xH(x,x) \vdash \exists x\exists y\neg(x=y)$

1		$\exists x\exists y(H(x,y) \vee H(y,x))$			
2		$\neg\exists xH(x,x)$			
		<hr/>			
3		u	v		$u = v$
					<hr/>
4					$H(u,v) \vee H(v,u)$
					<hr/>
5					$H(u,v)$
					<hr/>
6					$H(u,u)$ $=E, 3, 5$

Ex10. Build a proof of $\exists x\exists y(H(x,y) \vee H(y,x)), \neg\exists xH(x,x) \vdash \exists x\exists y\neg(x=y)$

1	$\exists x\exists y(H(x,y) \vee H(y,x))$				
2	$\neg\exists xH(x,x)$				
<hr/>					
3	u	v		$u = v$	
<hr/>					
4				$H(u,v) \vee H(v,u)$	
<hr/>					
5				$H(u,v)$	
<hr/>					
6				$H(u,u)$	$=E, 3, 5$
<hr/>					
7				$\exists xH(x,x)$	$\exists I, 5$

Ex10. Build a proof of $\exists x\exists y(H(x,y) \vee H(y,x)), \neg\exists xH(x,x) \vdash \exists x\exists y\neg(x=y)$

1	$\exists x\exists y(H(x,y) \vee H(y,x))$				
2	$\neg\exists xH(x,x)$				
3	u	v	$u = v$		
4			$H(u,v) \vee H(v,u)$		
5			$H(u,v)$		
6			$H(u,u)$	$=E, 3, 5$	
7			$\exists xH(x,x)$	$\exists I, 5$	
8			\perp	$\perp I, 2, 7$	

Ex10. Build a proof of $\exists x\exists y(H(x, y) \vee H(y, x)), \neg\exists xH(x, x) \vdash \exists x\exists y\neg(x = y)$

Ex10. Build a proof of $\exists x\exists y(H(x, y) \vee H(y, x)), \neg\exists xH(x, x) \vdash \exists x\exists y\neg(x = y)$

8 | | | | | \vdots

Ex10. Build a proof of $\exists x\exists y(H(x, y) \vee H(y, x)), \neg\exists xH(x, x) \vdash \exists x\exists y\neg(x = y)$

8						⋮
9						<u>$H(v, u)$</u>

Ex10. Build a proof of $\exists x\exists y(H(x, y) \vee H(y, x)), \neg\exists xH(x, x) \vdash \exists x\exists y\neg(x = y)$

8						⋮		
9							$H(v, u)$	
10							$H(v, v)$	=E, 3, 9

Ex10. Build a proof of $\exists x\exists y(H(x, y) \vee H(y, x)), \neg\exists xH(x, x) \vdash \exists x\exists y\neg(x = y)$

8						⋮		
9							$H(v, u)$	
10							$H(v, v)$	=E, 3, 9
11							$\exists xH(x, x)$	\exists I, 10

Ex10. Build a proof of $\exists x\exists y(H(x, y) \vee H(y, x)), \neg\exists xH(x, x) \vdash \exists x\exists y\neg(x = y)$

8						⋮		
9							$H(v, u)$	
10							$H(v, v)$	=E, 3, 9
11							$\exists xH(x, x)$	$\exists I, 10$
12							\perp	$\perp I, 2, 7$

Ex10. Build a proof of $\exists x\exists y(H(x, y) \vee H(y, x)), \neg\exists xH(x, x) \vdash \exists x\exists y\neg(x = y)$

8				⋮		
9					$H(v, u)$	
10					$H(v, v)$	=E, 3, 9
11					$\exists xH(x, x)$	$\exists I, 10$
12					\perp	$\perp I, 2, 7$
13				\perp		$\forall E, 4, 5-8, 9-12$

Ex10. Build a proof of $\exists x\exists y(H(x, y) \vee H(y, x)), \neg\exists xH(x, x) \vdash \exists x\exists y\neg(x = y)$

8					⋮			
9							$H(v, u)$	
10							$H(v, v)$	=E, 3, 9
11							$\exists xH(x, x)$	$\exists I, 10$
12							\perp	$\perp I, 2, 7$
13					\perp			$\forall E, 4, 5-8, 9-12$
14				\perp				$\exists E, 1, 4-13$

Ex10. Build a proof of $\exists x\exists y(H(x, y) \vee H(y, x)), \neg\exists xH(x, x) \vdash \exists x\exists y\neg(x = y)$

8				⋮		
9						$H(v, u)$
10						$H(v, v)$ =E, 3, 9
11						$\exists xH(x, x)$ \exists I, 10
12						\perp \perp I, 2, 7
13					\perp	\forall E, 4, 5–8, 9–12
14					\perp	\exists E, 1, 4–13
15						$\neg(u = v)$ \neg I, 3–14

Ex10. Build a proof of $\exists x\exists y(H(x, y) \vee H(y, x)), \neg\exists xH(x, x) \vdash \exists x\exists y\neg(x = y)$

8				⋮		
9						$H(v, u)$
10						$H(v, v)$ =E, 3, 9
11						$\exists xH(x, x)$ \exists I, 10
12						\perp \perp I, 2, 7
13					⊥	\forall E, 4, 5–8, 9–12
14					⊥	\exists E, 1, 4–13
15						$\neg(u = v)$ \neg I, 3–14
16						$\exists y\neg(u = y)$ \exists I, 15

Ex10. Build a proof of $\exists x\exists y(H(x, y) \vee H(y, x)), \neg\exists xH(x, x) \vdash \exists x\exists y\neg(x = y)$

8				⋮			
9						$H(v, u)$	
10						$H(v, v)$	=E, 3, 9
11						$\exists xH(x, x)$	$\exists I, 10$
12						\perp	$\perp I, 2, 7$
13				\perp			$\forall E, 4, 5-8, 9-12$
14			\perp				$\exists E, 1, 4-13$
15		$\neg(u = v)$					$\neg I, 3-14$
16		$\exists y\neg(u = y)$					$\exists I, 15$
17	$\exists x\exists y\neg(x = y)$						$\exists I, 16$

Ex11. Build a proof of $\exists y\exists xQ(y, x) \vdash \exists x\exists yQ(y, x)$

Ex11. Build a proof of $\exists y\exists xQ(y, x) \vdash \exists x\exists yQ(y, x)$

1 $\exists y\exists xQ(y, x)$

Ex11. Build a proof of $\exists y\exists xQ(y, x) \vdash \exists x\exists yQ(y, x)$

$$\begin{array}{l|l} 1 & \exists y\exists xQ(y, x) \\ \hline 2 & u \mid \exists xQ(u, x) \end{array}$$

Ex11. Build a proof of $\exists y\exists xQ(y, x) \vdash \exists x\exists yQ(y, x)$

1		$\exists y\exists xQ(y, x)$

2		u $\exists xQ(u, x)$

3		v $Q(u, v)$

Ex11. Build a proof of $\exists y\exists xQ(y, x) \vdash \exists x\exists yQ(y, x)$

1		$\exists y\exists xQ(y, x)$			

2		u $\exists xQ(u, x)$			

3			v $Q(u, v)$		

4				$\exists yQ(y, v)$	$\exists I, 3$

Ex11. Build a proof of $\exists y\exists xQ(y, x) \vdash \exists x\exists yQ(y, x)$

1		$\exists y\exists xQ(y, x)$	
<hr/>			
2		u $\exists xQ(u, x)$	
<hr/>			
3		v $Q(u, v)$	
<hr/>			
4		$\exists yQ(y, v)$	$\exists I, 3$
<hr/>			
5		$\exists yQ(y, v)$	$\exists E, 2, 3-4$

Ex11. Build a proof of $\exists y\exists xQ(y, x) \vdash \exists x\exists yQ(y, x)$

1		$\exists y\exists xQ(y, x)$	
<hr/>			
2	u	$\exists xQ(u, x)$	
<hr/>			
3	v	$Q(u, v)$	
<hr/>			
4		$\exists yQ(y, v)$	$\exists I, 3$
5		$\exists yQ(y, v)$	$\exists E, 2, 3-4$
6		$\exists x\exists yQ(y, x)$	$\exists I, 5$

Ex11. Build a proof of $\exists y\exists xQ(y, x) \vdash \exists x\exists yQ(y, x)$

1		$\exists y\exists xQ(y, x)$	
<hr/>			
2		u $\exists xQ(u, x)$	
<hr/>			
3		v $Q(u, v)$	
<hr/>			
4		$\exists yQ(y, v)$	$\exists I, 3$
5		$\exists yQ(y, v)$	$\exists E, 2, 3-4$
6		$\exists x\exists yQ(y, x)$	$\exists I, 5$
7		$\exists x\exists yQ(y, x)$	$\exists E, 1, 2-6$

Ex12. Build a proof of $\vdash (\exists xP(x) \rightarrow \forall xR(x)) \rightarrow (\forall x(P(x) \rightarrow Q(x)))$

Ex12. Build a proof of $\vdash (\exists xP(x) \rightarrow \forall xR(x)) \rightarrow (\forall x(P(x) \rightarrow Q(x)))$

1	$\exists xP(x) \rightarrow \forall xR(x)$
2	u $P(u)$

Ex12. Build a proof of $\vdash (\exists xP(x) \rightarrow \forall xR(x)) \rightarrow (\forall x(P(x) \rightarrow Q(x)))$

1			$\exists xP(x) \rightarrow \forall xR(x)$	
2		u	$P(u)$	
3			$\exists xP(x)$	$\exists I, 2$

Ex12. Build a proof of $\vdash (\exists xP(x) \rightarrow \forall xR(x)) \rightarrow (\forall x(P(x) \rightarrow Q(x)))$

1			$\exists xP(x) \rightarrow \forall xR(x)$	
2			u $P(u)$	
3			$\exists xP(x)$	$\exists I, 2$
4			$\forall xR(x)$	$\Rightarrow E, 1, 3$

Ex12. Build a proof of $\vdash (\exists xP(x) \rightarrow \forall xR(x)) \rightarrow (\forall x(P(x) \rightarrow Q(x)))$

1			$\exists xP(x) \rightarrow \forall xR(x)$	
2		u	$P(u)$	
3			$\exists xP(x)$	$\exists I, 2$
4			$\forall xR(x)$	$\Rightarrow E, 1, 3$
5			$R(u)$	$\forall E, 4$

Ex12. Build a proof of $\vdash (\exists xP(x) \rightarrow \forall xR(x)) \rightarrow (\forall x(P(x) \rightarrow Q(x)))$

1			$\exists xP(x) \rightarrow \forall xR(x)$	
2			u $P(u)$	
3			$\exists xP(x)$	$\exists I, 2$
4			$\forall xR(x)$	$\Rightarrow E, 1, 3$
5			$R(u)$	$\forall E, 4$
6			$P(u) \rightarrow Q(u)$	$\Rightarrow I, 2-5$

Ex12. Build a proof of $\vdash (\exists xP(x) \rightarrow \forall xR(x)) \rightarrow (\forall x(P(x) \rightarrow Q(x)))$

1			$\exists xP(x) \rightarrow \forall xR(x)$	
2			u $P(u)$	
3			$\exists xP(x)$	$\exists I, 2$
4			$\forall xR(x)$	$\Rightarrow E, 1, 3$
5			$R(u)$	$\forall E, 4$
6			$P(u) \rightarrow Q(u)$	$\Rightarrow I, 2-5$
7			$\forall x(P(x) \rightarrow Q(x))$	$\forall I, 2-6$

Ex12. Build a proof of $\vdash (\exists xP(x) \rightarrow \forall xR(x)) \rightarrow (\forall x(P(x) \rightarrow Q(x)))$

1	$\exists xP(x) \rightarrow \forall xR(x)$ <hr style="border: 0.5px solid black;"/>	
2	u $P(u)$ <hr style="border: 0.5px solid black;"/>	
3	$\exists xP(x)$	$\exists I, 2$
4	$\forall xR(x)$	$\Rightarrow E, 1, 3$
5	$R(u)$	$\forall E, 4$
6	$P(u) \rightarrow Q(u)$	$\Rightarrow I, 2-5$
7	$\forall x(P(x) \rightarrow Q(x))$	$\forall I, 2-6$
8	$(\exists xP(x) \rightarrow \forall xR(x)) \rightarrow (\forall x(P(x) \rightarrow Q(x)))$	$\Rightarrow I, 1-7$

Ex13. Build a proof of $\vdash x = f(y) \rightarrow \forall z(P(x, z) \rightarrow P(f(y), z))$

Ex13. Build a proof of $\vdash x = f(y) \rightarrow \forall z(P(x, z) \rightarrow P(f(y), z))$

1 | $x = f(y)$

Ex13. Build a proof of $\vdash x = f(y) \rightarrow \forall z(P(x, z) \rightarrow P(f(y), z))$

1			$x = f(y)$
		<hr/>	
2			$P(x, u)$
		<hr/>	

Ex13. Build a proof of $\vdash x = f(y) \rightarrow \forall z(P(x, z) \rightarrow P(f(y), z))$

1			$x = f(y)$	
2			u $P(x, u)$	
3			$P(f(y), u)$	$=E, 1, 2$

Ex13. Build a proof of $\vdash x = f(y) \rightarrow \forall z(P(x, z) \rightarrow P(f(y), z))$

1			$x = f(y)$	
			<hr/>	
2			u	
				$P(x, u)$
				<hr/>
3				$P(f(y), u)$
				$\Rightarrow E, 1, 2$
4				$P(x, u) \rightarrow P(f(y), u)$
				$\Rightarrow I, 2-3$

Ex13. Build a proof of $\vdash x = f(y) \rightarrow \forall z(P(x, z) \rightarrow P(f(y), z))$

1			$x = f(y)$		

2			u		
				$P(x, u)$	

3				$P(f(y), u)$	$=E, 1, 2$

4				$P(x, u) \rightarrow P(f(y), u)$	$\Rightarrow I, 2-3$
5				$\forall z(P(x, z) \rightarrow P(f(y), z))$	$\forall I, 2-4$

Ex13. Build a proof of $\vdash x = f(y) \rightarrow \forall z(P(x, z) \rightarrow P(f(y), z))$

1			$x = f(y)$		

2			u		
				$P(x, u)$	

3				$P(f(y), u)$	$=E, 1, 2$

4				$P(x, u) \rightarrow P(f(y), u)$	$\Rightarrow I, 2-3$

5				$\forall z(P(x, z) \rightarrow P(f(y), z))$	$\forall I, 2-4$
6				$x = f(y) \rightarrow \forall z(P(x, z) \rightarrow P(f(y), z))$	$\Rightarrow I, 1-5$

Ex14. Build a proof of $\forall x \forall y \forall u \forall v (x = u \rightarrow (y = v \rightarrow f(x, y) = f(u, v)))$

Ex14. Build a proof of $\forall x \forall y \forall u \forall v (x = u \rightarrow (y = v \rightarrow f(x, y) = f(u, v)))$

1 | a | b | c | d | $a = b$

Ex14. Build a proof of $\forall x \forall y \forall u \forall v (x = u \rightarrow (y = v \rightarrow f(x, y) = f(u, v)))$

1		<i>a</i>		<i>b</i>		<i>c</i>		<i>d</i>				<i>a = b</i>		
2														<i>c = d</i>

Ex14. Build a proof of $\forall x \forall y \forall u \forall v (x = u \rightarrow (y = v \rightarrow f(x, y) = f(u, v)))$

1		<i>a</i>		<i>b</i>		<i>c</i>		<i>d</i>			<i>a = b</i>		
2												<i>c = d</i>	
3												<i>f(a, c) = f(a, c)</i>	=I

Ex14. Build a proof of $\forall x \forall y \forall u \forall v (x = u \rightarrow (y = v \rightarrow f(x, y) = f(u, v)))$

1		a		b		c		d			a = b		
2												c = d	
3												f(a, c) = f(a, c)	=I
4												f(a, c) = f(b, c)	=E, 1, 3

Ex14. Build a proof of $\forall x \forall y \forall u \forall v (x = u \rightarrow (y = v \rightarrow f(x, y) = f(u, v)))$

1		a		b		c		d			a = b		
2												c = d	
3												f(a, c) = f(a, c)	=I
4												f(a, c) = f(b, c)	=E, 1, 3
5												f(a, c) = f(b, d)	=E, 2, 4

Ex14. Build a proof of $\forall x \forall y \forall u \forall v (x = u \rightarrow (y = v \rightarrow f(x, y) = f(u, v)))$

1		a		b		c		d		a = b	
2										c = d	
3										f(a, c) = f(a, c)	=I
4										f(a, c) = f(b, c)	=E, 1, 3
5										f(a, c) = f(b, d)	=E, 2, 4
6										c = d \rightarrow f(a, c) = f(b, d)	\Rightarrow I, 2-5

Ex14. Build a proof of $\forall x \forall y \forall u \forall v (x = u \rightarrow (y = v \rightarrow f(x, y) = f(u, v)))$

1	a	b	c	d	a = b		
2					c = d		
3					f(a, c) = f(a, c)	=I	
4					f(a, c) = f(b, c)	=E, 1, 3	
5					f(a, c) = f(b, d)	=E, 2, 4	
6					c = d → f(a, c) = f(b, d)	⇒I, 2-5	
7					a = b → (c = d → f(a, c) = f(b, d))	⇒I, 1-6	

Ex14. Build a proof of $\forall x \forall y \forall u \forall v (x = u \rightarrow (y = v \rightarrow f(x, y) = f(u, v)))$

1		a	b	c	d		$a = b$	
2							$c = d$	
3							$f(a, c) = f(a, c)$	=I
4							$f(a, c) = f(b, c)$	=E, 1, 3
5							$f(a, c) = f(b, d)$	=E, 2, 4
6							$c = d \rightarrow f(a, c) = f(b, d)$	\Rightarrow I, 2-5
7							$a = b \rightarrow (c = d \rightarrow f(a, c) = f(b, d))$	\Rightarrow I, 1-6
8							$\forall v (a = b \rightarrow (c = v \rightarrow f(a, c) = f(b, v)))$	\forall I, 7

Ex14. Build a proof of $\forall x \forall y \forall u \forall v (x = u \rightarrow (y = v \rightarrow f(x, y) = f(u, v)))$

1		a	b	c	d	$a = b$	
2						$c = d$	
3						$f(a, c) = f(a, c)$	=I
4						$f(a, c) = f(b, c)$	=E, 1, 3
5						$f(a, c) = f(b, d)$	=E, 2, 4
6						$c = d \rightarrow f(a, c) = f(b, d)$	\Rightarrow I, 2-5
7						$a = b \rightarrow (c = d \rightarrow f(a, c) = f(b, d))$	\Rightarrow I, 1-6
8						$\forall v (a = b \rightarrow (c = v \rightarrow f(a, c) = f(b, v)))$	\forall I, 7
9						$\forall u \forall v (a = u \rightarrow (c = v \rightarrow f(a, c) = f(u, v)))$	\forall I, 8

Ex14. Build a proof of $\forall x \forall y \forall u \forall v (x = u \rightarrow (y = v \rightarrow f(x, y) = f(u, v)))$

1		a	b	c	d	$a = b$	
2						$c = d$	
3						$f(a, c) = f(a, c)$	=I
4						$f(a, c) = f(b, c)$	=E, 1, 3
5						$f(a, c) = f(b, d)$	=E, 2, 4
6						$c = d \rightarrow f(a, c) = f(b, d)$	\Rightarrow I, 2-5
7						$a = b \rightarrow (c = d \rightarrow f(a, c) = f(b, d))$	\Rightarrow I, 1-6
8						$\forall v (a = b \rightarrow (c = v \rightarrow f(a, c) = f(b, v)))$	\forall I, 7
9						$\forall u \forall v (a = u \rightarrow (c = v \rightarrow f(a, c) = f(u, v)))$	\forall I, 8
10						$\forall y \forall u \forall v (a = u \rightarrow (y = v \rightarrow f(a, y) = f(u, v)))$	\forall I, 9

Ex14. Build a proof of $\forall x \forall y \forall u \forall v (x = u \rightarrow (y = v \rightarrow f(x, y) = f(u, v)))$

1	a	b	c	d	a = b	
2					c = d	
3					f(a, c) = f(a, c)	=I
4					f(a, c) = f(b, c)	=E, 1, 3
5					f(a, c) = f(b, d)	=E, 2, 4
6					c = d → f(a, c) = f(b, d)	⇒I, 2-5
7					a = b → (c = d → f(a, c) = f(b, d))	⇒I, 1-6
8					∀v (a = b → (c = v → f(a, c) = f(b, v)))	∀I, 7
9					∀u ∀v (a = u → (c = v → f(a, c) = f(u, v)))	∀I, 8
10					∀y ∀u ∀v (a = u → (y = v → f(a, y) = f(u, v)))	∀I, 9
11					∀x ∀y ∀u ∀v (x = u → (y = v → f(x, y) = f(u, v)))	∀I, 10

Ex15. Build a proof of $\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x)) \vdash \forall x\forall y((P(x) \wedge P(y)) \rightarrow x = y)$

Ex15. Build a proof of $\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x)) \vdash \forall x\forall y((P(x) \wedge P(y)) \rightarrow x = y)$

1 $\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x))$

Ex15. Build a proof of $\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x)) \vdash \forall x\forall y((P(x) \wedge P(y)) \rightarrow x = y)$

$$\begin{array}{l} 1 \quad | \quad \exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x)) \\ 2 \quad | \quad a \quad | \quad b \quad | \quad \underline{P(a) \wedge P(b)} \end{array}$$

Ex15. Build a proof of $\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x)) \vdash \forall x\forall y((P(x) \wedge P(y)) \rightarrow x = y)$

1		$\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x))$	
2		a	b $P(a) \wedge P(b)$
3			$P(a) \wedge (P(b) \rightarrow b = a)$

Ex15. Build a proof of $\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x)) \vdash \forall x\forall y((P(x) \wedge P(y)) \rightarrow x = y)$

1		$\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x))$		
2		a	b	$P(a) \wedge P(b)$
3				$P(a) \wedge (P(b) \rightarrow b = a)$
4				$P(b)$ $\wedge E, 2$

Ex15. Build a proof of $\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x)) \vdash \forall x\forall y((P(x) \wedge P(y)) \rightarrow x = y)$

1	$\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x))$			
2	a	b	$P(a) \wedge P(b)$	
3			$P(a) \wedge (P(b) \rightarrow b = a)$	
4			$P(b)$	$\wedge E, 2$
5			$P(b) \rightarrow b = a$	$\wedge E, 3$

Ex15. Build a proof of $\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x)) \vdash \forall x\forall y((P(x) \wedge P(y)) \rightarrow x = y)$

1	$\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x))$			
2	a	b	$P(a) \wedge P(b)$	
3			$P(a) \wedge (P(b) \rightarrow b = a)$	
4			$P(b)$	$\wedge E, 2$
5			$P(b) \rightarrow b = a$	$\wedge E, 3$
6			$b = a$	$\Rightarrow E, 4, 5$

Ex15. Build a proof of $\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x)) \vdash \forall x\forall y((P(x) \wedge P(y)) \rightarrow x = y)$

1	$\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x))$								
2	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;">a</td> <td style="border-right: 1px solid black; padding-right: 5px;">b</td> <td style="padding-left: 5px;"> $P(a) \wedge P(b)$ </td> </tr> </table>	a	b	$P(a) \wedge P(b)$					
a	b	$P(a) \wedge P(b)$							
3	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(a) \wedge (P(b) \rightarrow b = a)$ </td> </tr> </table> </td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(a) \wedge (P(b) \rightarrow b = a)$ </td> </tr> </table>		$P(a) \wedge (P(b) \rightarrow b = a)$			
		<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(a) \wedge (P(b) \rightarrow b = a)$ </td> </tr> </table>		$P(a) \wedge (P(b) \rightarrow b = a)$					
	$P(a) \wedge (P(b) \rightarrow b = a)$								
4	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(b)$ </td> </tr> </table> </td> <td style="padding-left: 10px;">$\wedge E, 2$</td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(b)$ </td> </tr> </table>		$P(b)$	$\wedge E, 2$		
		<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(b)$ </td> </tr> </table>		$P(b)$	$\wedge E, 2$				
	$P(b)$								
5	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(b) \rightarrow b = a$ </td> </tr> </table> </td> <td style="padding-left: 10px;">$\wedge E, 3$</td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(b) \rightarrow b = a$ </td> </tr> </table>		$P(b) \rightarrow b = a$	$\wedge E, 3$		
		<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(b) \rightarrow b = a$ </td> </tr> </table>		$P(b) \rightarrow b = a$	$\wedge E, 3$				
	$P(b) \rightarrow b = a$								
6	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $b = a$ </td> </tr> </table> </td> <td style="padding-left: 10px;">$\Rightarrow E, 4, 5$</td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $b = a$ </td> </tr> </table>		$b = a$	$\Rightarrow E, 4, 5$		
		<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $b = a$ </td> </tr> </table>		$b = a$	$\Rightarrow E, 4, 5$				
	$b = a$								
7	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $b = b$ </td> </tr> </table> </td> <td style="padding-left: 10px;">$=I$</td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $b = b$ </td> </tr> </table>		$b = b$	$=I$		
		<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $b = b$ </td> </tr> </table>		$b = b$	$=I$				
	$b = b$								

Ex15. Build a proof of $\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x)) \vdash \forall x\forall y((P(x) \wedge P(y)) \rightarrow x = y)$

1	$\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x))$							
2	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;">a</td> <td style="border-right: 1px solid black; padding-right: 5px;">b</td> <td style="padding-left: 10px;"> $P(a) \wedge P(b)$ </td> </tr> </table>	a	b	$P(a) \wedge P(b)$				
a	b	$P(a) \wedge P(b)$						
3	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 10px;"> <table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $P(a) \wedge (P(b) \rightarrow b = a)$ </td> </tr> </table> </td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $P(a) \wedge (P(b) \rightarrow b = a)$ </td> </tr> </table>	$P(a) \wedge (P(b) \rightarrow b = a)$			
		<table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $P(a) \wedge (P(b) \rightarrow b = a)$ </td> </tr> </table>	$P(a) \wedge (P(b) \rightarrow b = a)$					
$P(a) \wedge (P(b) \rightarrow b = a)$								
4	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 10px;"> <table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $P(b)$ </td> <td style="padding-left: 10px;">$\wedge E, 2$</td> </tr> </table> </td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $P(b)$ </td> <td style="padding-left: 10px;">$\wedge E, 2$</td> </tr> </table>	$P(b)$	$\wedge E, 2$		
		<table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $P(b)$ </td> <td style="padding-left: 10px;">$\wedge E, 2$</td> </tr> </table>	$P(b)$	$\wedge E, 2$				
$P(b)$	$\wedge E, 2$							
5	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 10px;"> <table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $P(b) \rightarrow b = a$ </td> <td style="padding-left: 10px;">$\wedge E, 3$</td> </tr> </table> </td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $P(b) \rightarrow b = a$ </td> <td style="padding-left: 10px;">$\wedge E, 3$</td> </tr> </table>	$P(b) \rightarrow b = a$	$\wedge E, 3$		
		<table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $P(b) \rightarrow b = a$ </td> <td style="padding-left: 10px;">$\wedge E, 3$</td> </tr> </table>	$P(b) \rightarrow b = a$	$\wedge E, 3$				
$P(b) \rightarrow b = a$	$\wedge E, 3$							
6	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 10px;"> <table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $b = a$ </td> <td style="padding-left: 10px;">$\Rightarrow E, 4, 5$</td> </tr> </table> </td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $b = a$ </td> <td style="padding-left: 10px;">$\Rightarrow E, 4, 5$</td> </tr> </table>	$b = a$	$\Rightarrow E, 4, 5$		
		<table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $b = a$ </td> <td style="padding-left: 10px;">$\Rightarrow E, 4, 5$</td> </tr> </table>	$b = a$	$\Rightarrow E, 4, 5$				
$b = a$	$\Rightarrow E, 4, 5$							
7	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 10px;"> <table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $b = b$ </td> <td style="padding-left: 10px;">$=I$</td> </tr> </table> </td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $b = b$ </td> <td style="padding-left: 10px;">$=I$</td> </tr> </table>	$b = b$	$=I$		
		<table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $b = b$ </td> <td style="padding-left: 10px;">$=I$</td> </tr> </table>	$b = b$	$=I$				
$b = b$	$=I$							
8	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 10px;"> <table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $a = b$ </td> <td style="padding-left: 10px;">$=E, 6, 7$</td> </tr> </table> </td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $a = b$ </td> <td style="padding-left: 10px;">$=E, 6, 7$</td> </tr> </table>	$a = b$	$=E, 6, 7$		
		<table style="border-collapse: collapse;"> <tr> <td style="padding-left: 10px;"> $a = b$ </td> <td style="padding-left: 10px;">$=E, 6, 7$</td> </tr> </table>	$a = b$	$=E, 6, 7$				
$a = b$	$=E, 6, 7$							

Ex15. Build a proof of $\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x)) \vdash \forall x\forall y((P(x) \wedge P(y)) \rightarrow x = y)$

1	$\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x))$								
2	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px; text-align: center;">a</td> <td style="border-right: 1px solid black; padding-right: 5px; text-align: center;">b</td> <td style="padding-left: 5px;"> $P(a) \wedge P(b)$ </td> </tr> </table>	a	b	$P(a) \wedge P(b)$					
a	b	$P(a) \wedge P(b)$							
3	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(a) \wedge (P(b) \rightarrow b = a)$ </td> </tr> </table> </td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(a) \wedge (P(b) \rightarrow b = a)$ </td> </tr> </table>		$P(a) \wedge (P(b) \rightarrow b = a)$			
		<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(a) \wedge (P(b) \rightarrow b = a)$ </td> </tr> </table>		$P(a) \wedge (P(b) \rightarrow b = a)$					
	$P(a) \wedge (P(b) \rightarrow b = a)$								
4	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(b)$ </td> <td style="padding-left: 10px; vertical-align: middle;">$\wedge E, 2$</td> </tr> </table> </td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(b)$ </td> <td style="padding-left: 10px; vertical-align: middle;">$\wedge E, 2$</td> </tr> </table>		$P(b)$	$\wedge E, 2$		
		<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(b)$ </td> <td style="padding-left: 10px; vertical-align: middle;">$\wedge E, 2$</td> </tr> </table>		$P(b)$	$\wedge E, 2$				
	$P(b)$	$\wedge E, 2$							
5	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(b) \rightarrow b = a$ </td> <td style="padding-left: 10px; vertical-align: middle;">$\wedge E, 3$</td> </tr> </table> </td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(b) \rightarrow b = a$ </td> <td style="padding-left: 10px; vertical-align: middle;">$\wedge E, 3$</td> </tr> </table>		$P(b) \rightarrow b = a$	$\wedge E, 3$		
		<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $P(b) \rightarrow b = a$ </td> <td style="padding-left: 10px; vertical-align: middle;">$\wedge E, 3$</td> </tr> </table>		$P(b) \rightarrow b = a$	$\wedge E, 3$				
	$P(b) \rightarrow b = a$	$\wedge E, 3$							
6	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $b = a$ </td> <td style="padding-left: 10px; vertical-align: middle;">$\Rightarrow E, 4, 5$</td> </tr> </table> </td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $b = a$ </td> <td style="padding-left: 10px; vertical-align: middle;">$\Rightarrow E, 4, 5$</td> </tr> </table>		$b = a$	$\Rightarrow E, 4, 5$		
		<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $b = a$ </td> <td style="padding-left: 10px; vertical-align: middle;">$\Rightarrow E, 4, 5$</td> </tr> </table>		$b = a$	$\Rightarrow E, 4, 5$				
	$b = a$	$\Rightarrow E, 4, 5$							
7	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $b = b$ </td> <td style="padding-left: 10px; vertical-align: middle;">$=I$</td> </tr> </table> </td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $b = b$ </td> <td style="padding-left: 10px; vertical-align: middle;">$=I$</td> </tr> </table>		$b = b$	$=I$		
		<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $b = b$ </td> <td style="padding-left: 10px; vertical-align: middle;">$=I$</td> </tr> </table>		$b = b$	$=I$				
	$b = b$	$=I$							
8	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $a = b$ </td> <td style="padding-left: 10px; vertical-align: middle;">$=E, 6, 7$</td> </tr> </table> </td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $a = b$ </td> <td style="padding-left: 10px; vertical-align: middle;">$=E, 6, 7$</td> </tr> </table>		$a = b$	$=E, 6, 7$		
		<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $a = b$ </td> <td style="padding-left: 10px; vertical-align: middle;">$=E, 6, 7$</td> </tr> </table>		$a = b$	$=E, 6, 7$				
	$a = b$	$=E, 6, 7$							
9	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $a = b$ </td> <td style="padding-left: 10px; vertical-align: middle;">$\exists E, 1, 3-8$</td> </tr> </table> </td> </tr> </table>			<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $a = b$ </td> <td style="padding-left: 10px; vertical-align: middle;">$\exists E, 1, 3-8$</td> </tr> </table>		$a = b$	$\exists E, 1, 3-8$		
		<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"></td> <td style="padding-left: 5px;"> $a = b$ </td> <td style="padding-left: 10px; vertical-align: middle;">$\exists E, 1, 3-8$</td> </tr> </table>		$a = b$	$\exists E, 1, 3-8$				
	$a = b$	$\exists E, 1, 3-8$							

Ex15. Build a proof of $\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x)) \vdash \forall x\forall y((P(x) \wedge P(y)) \rightarrow x = y)$

1	$\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x))$																																							
2	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px; width: 5%;">a</td> <td style="border-right: 1px solid black; padding-right: 5px; width: 5%;">b</td> <td style="padding-left: 5px;">$P(a) \wedge P(b)$</td> </tr> </table>	a	b	$P(a) \wedge P(b)$																																				
a	b	$P(a) \wedge P(b)$																																						
3	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="padding-left: 5px;">$P(a) \wedge (P(b) \rightarrow b = a)$</td> </tr> </table> </td> <td style="padding-left: 5px;">$P(b)$</td> <td style="padding-left: 5px;">$\wedge E, 2$</td> </tr> <tr> <td style="padding-left: 5px;">$P(b) \rightarrow b = a$</td> <td></td> <td style="padding-left: 5px;">$\wedge E, 3$</td> </tr> <tr> <td style="padding-left: 5px;">$b = a$</td> <td></td> <td style="padding-left: 5px;">$\Rightarrow E, 4, 5$</td> </tr> <tr> <td style="padding-left: 5px;">$b = b$</td> <td></td> <td style="padding-left: 5px;">$=I$</td> </tr> <tr> <td style="padding-left: 5px;">$a = b$</td> <td></td> <td style="padding-left: 5px;">$=E, 6, 7$</td> </tr> </table> </td> <td></td> </tr> <tr> <td style="text-align: right; vertical-align: top;">4</td> <td style="border-left: 1px solid black; padding-left: 5px;">$P(b)$</td> <td style="padding-left: 5px;">$\wedge E, 2$</td> </tr> <tr> <td style="text-align: right; vertical-align: top;">5</td> <td style="border-left: 1px solid black; padding-left: 5px;">$P(b) \rightarrow b = a$</td> <td style="padding-left: 5px;">$\wedge E, 3$</td> </tr> <tr> <td style="text-align: right; vertical-align: top;">6</td> <td style="border-left: 1px solid black; padding-left: 5px;">$b = a$</td> <td style="padding-left: 5px;">$\Rightarrow E, 4, 5$</td> </tr> <tr> <td style="text-align: right; vertical-align: top;">7</td> <td style="border-left: 1px solid black; padding-left: 5px;">$b = b$</td> <td style="padding-left: 5px;">$=I$</td> </tr> <tr> <td style="text-align: right; vertical-align: top;">8</td> <td style="border-left: 1px solid black; padding-left: 5px;">$a = b$</td> <td style="padding-left: 5px;">$=E, 6, 7$</td> </tr> <tr> <td style="text-align: right; vertical-align: top;">9</td> <td style="border-left: 1px solid black; padding-left: 5px;">$a = b$</td> <td style="padding-left: 5px;">$\exists E, 1, 3-8$</td> </tr> <tr> <td style="text-align: right; vertical-align: top;">10</td> <td style="border-left: 1px solid black; padding-left: 5px;">$(P(a) \wedge P(b)) \rightarrow a = b$</td> <td style="padding-left: 5px;">$\Rightarrow I, 2-9$</td> </tr> </table>	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="padding-left: 5px;">$P(a) \wedge (P(b) \rightarrow b = a)$</td> </tr> </table> </td> <td style="padding-left: 5px;">$P(b)$</td> <td style="padding-left: 5px;">$\wedge E, 2$</td> </tr> <tr> <td style="padding-left: 5px;">$P(b) \rightarrow b = a$</td> <td></td> <td style="padding-left: 5px;">$\wedge E, 3$</td> </tr> <tr> <td style="padding-left: 5px;">$b = a$</td> <td></td> <td style="padding-left: 5px;">$\Rightarrow E, 4, 5$</td> </tr> <tr> <td style="padding-left: 5px;">$b = b$</td> <td></td> <td style="padding-left: 5px;">$=I$</td> </tr> <tr> <td style="padding-left: 5px;">$a = b$</td> <td></td> <td style="padding-left: 5px;">$=E, 6, 7$</td> </tr> </table>	<table style="border-collapse: collapse;"> <tr> <td style="padding-left: 5px;">$P(a) \wedge (P(b) \rightarrow b = a)$</td> </tr> </table>	$P(a) \wedge (P(b) \rightarrow b = a)$	$P(b)$	$\wedge E, 2$	$P(b) \rightarrow b = a$		$\wedge E, 3$	$b = a$		$\Rightarrow E, 4, 5$	$b = b$		$=I$	$a = b$		$=E, 6, 7$		4	$P(b)$	$\wedge E, 2$	5	$P(b) \rightarrow b = a$	$\wedge E, 3$	6	$b = a$	$\Rightarrow E, 4, 5$	7	$b = b$	$=I$	8	$a = b$	$=E, 6, 7$	9	$a = b$	$\exists E, 1, 3-8$	10	$(P(a) \wedge P(b)) \rightarrow a = b$	$\Rightarrow I, 2-9$
<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;"> <table style="border-collapse: collapse;"> <tr> <td style="padding-left: 5px;">$P(a) \wedge (P(b) \rightarrow b = a)$</td> </tr> </table> </td> <td style="padding-left: 5px;">$P(b)$</td> <td style="padding-left: 5px;">$\wedge E, 2$</td> </tr> <tr> <td style="padding-left: 5px;">$P(b) \rightarrow b = a$</td> <td></td> <td style="padding-left: 5px;">$\wedge E, 3$</td> </tr> <tr> <td style="padding-left: 5px;">$b = a$</td> <td></td> <td style="padding-left: 5px;">$\Rightarrow E, 4, 5$</td> </tr> <tr> <td style="padding-left: 5px;">$b = b$</td> <td></td> <td style="padding-left: 5px;">$=I$</td> </tr> <tr> <td style="padding-left: 5px;">$a = b$</td> <td></td> <td style="padding-left: 5px;">$=E, 6, 7$</td> </tr> </table>	<table style="border-collapse: collapse;"> <tr> <td style="padding-left: 5px;">$P(a) \wedge (P(b) \rightarrow b = a)$</td> </tr> </table>	$P(a) \wedge (P(b) \rightarrow b = a)$	$P(b)$	$\wedge E, 2$	$P(b) \rightarrow b = a$		$\wedge E, 3$	$b = a$		$\Rightarrow E, 4, 5$	$b = b$		$=I$	$a = b$		$=E, 6, 7$																								
<table style="border-collapse: collapse;"> <tr> <td style="padding-left: 5px;">$P(a) \wedge (P(b) \rightarrow b = a)$</td> </tr> </table>	$P(a) \wedge (P(b) \rightarrow b = a)$	$P(b)$	$\wedge E, 2$																																					
$P(a) \wedge (P(b) \rightarrow b = a)$																																								
$P(b) \rightarrow b = a$		$\wedge E, 3$																																						
$b = a$		$\Rightarrow E, 4, 5$																																						
$b = b$		$=I$																																						
$a = b$		$=E, 6, 7$																																						
4	$P(b)$	$\wedge E, 2$																																						
5	$P(b) \rightarrow b = a$	$\wedge E, 3$																																						
6	$b = a$	$\Rightarrow E, 4, 5$																																						
7	$b = b$	$=I$																																						
8	$a = b$	$=E, 6, 7$																																						
9	$a = b$	$\exists E, 1, 3-8$																																						
10	$(P(a) \wedge P(b)) \rightarrow a = b$	$\Rightarrow I, 2-9$																																						

Ex15. Build a proof of $\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x)) \vdash \forall x\forall y((P(x) \wedge P(y)) \rightarrow x = y)$

1	$\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x))$			
2	a	b	$P(a) \wedge P(b)$	
3			$P(a) \wedge (P(b) \rightarrow b = a)$	
4			$P(b)$	$\wedge E, 2$
5			$P(b) \rightarrow b = a$	$\wedge E, 3$
6			$b = a$	$\Rightarrow E, 4, 5$
7			$b = b$	$=I$
8			$a = b$	$=E, 6, 7$
9			$a = b$	$\exists E, 1, 3-8$
10			$(P(a) \wedge P(b)) \rightarrow a = b$	$\Rightarrow I, 2-9$
11			$\forall y((P(a) \wedge P(y)) \rightarrow a = y)$	$\forall I, 10$

Ex15. Build a proof of $\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x)) \vdash \forall x\forall y((P(x) \wedge P(y)) \rightarrow x = y)$

1	$\exists x\forall y(P(x) \wedge (P(y) \rightarrow y = x))$					
2	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding: 5px; width: 20px;">a</td> <td style="border-right: 1px solid black; padding: 5px; width: 20px;">b</td> <td style="padding: 5px;">$P(a) \wedge P(b)$</td> </tr> </table>	a	b	$P(a) \wedge P(b)$		
a	b	$P(a) \wedge P(b)$				
3	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding: 5px; width: 20px;"></td> <td style="border-right: 1px solid black; padding: 5px; width: 20px;"></td> <td style="padding: 5px;">$P(a) \wedge (P(b) \rightarrow b = a)$</td> </tr> </table>			$P(a) \wedge (P(b) \rightarrow b = a)$		
		$P(a) \wedge (P(b) \rightarrow b = a)$				
4	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding: 5px; width: 20px;"></td> <td style="border-right: 1px solid black; padding: 5px; width: 20px;"></td> <td style="padding: 5px;">$P(b)$</td> <td style="padding: 5px;">$\wedge E, 2$</td> </tr> </table>			$P(b)$	$\wedge E, 2$	
		$P(b)$	$\wedge E, 2$			
5	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding: 5px; width: 20px;"></td> <td style="border-right: 1px solid black; padding: 5px; width: 20px;"></td> <td style="padding: 5px;">$P(b) \rightarrow b = a$</td> <td style="padding: 5px;">$\wedge E, 3$</td> </tr> </table>			$P(b) \rightarrow b = a$	$\wedge E, 3$	
		$P(b) \rightarrow b = a$	$\wedge E, 3$			
6	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding: 5px; width: 20px;"></td> <td style="border-right: 1px solid black; padding: 5px; width: 20px;"></td> <td style="padding: 5px;">$b = a$</td> <td style="padding: 5px;">$\Rightarrow E, 4, 5$</td> </tr> </table>			$b = a$	$\Rightarrow E, 4, 5$	
		$b = a$	$\Rightarrow E, 4, 5$			
7	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding: 5px; width: 20px;"></td> <td style="border-right: 1px solid black; padding: 5px; width: 20px;"></td> <td style="padding: 5px;">$b = b$</td> <td style="padding: 5px;">$=I$</td> </tr> </table>			$b = b$	$=I$	
		$b = b$	$=I$			
8	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding: 5px; width: 20px;"></td> <td style="border-right: 1px solid black; padding: 5px; width: 20px;"></td> <td style="padding: 5px;">$a = b$</td> <td style="padding: 5px;">$=E, 6, 7$</td> </tr> </table>			$a = b$	$=E, 6, 7$	
		$a = b$	$=E, 6, 7$			
9	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding: 5px; width: 20px;"></td> <td style="padding: 5px;">$a = b$</td> <td style="padding: 5px;">$\exists E, 1, 3-8$</td> </tr> </table>		$a = b$	$\exists E, 1, 3-8$		
	$a = b$	$\exists E, 1, 3-8$				
10	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding: 5px; width: 20px;"></td> <td style="padding: 5px;">$(P(a) \wedge P(b)) \rightarrow a = b$</td> <td style="padding: 5px;">$\Rightarrow I, 2-9$</td> </tr> </table>		$(P(a) \wedge P(b)) \rightarrow a = b$	$\Rightarrow I, 2-9$		
	$(P(a) \wedge P(b)) \rightarrow a = b$	$\Rightarrow I, 2-9$				
11	<table style="border-collapse: collapse;"> <tr> <td style="padding: 5px;">$\forall y((P(a) \wedge P(y)) \rightarrow a = y)$</td> <td style="padding: 5px;">$\forall I, 10$</td> </tr> </table>	$\forall y((P(a) \wedge P(y)) \rightarrow a = y)$	$\forall I, 10$			
$\forall y((P(a) \wedge P(y)) \rightarrow a = y)$	$\forall I, 10$					
12	$\forall x\forall y((P(x) \wedge P(y)) \rightarrow x = y)$	$\forall I, 11$				